

BARI-TH 444/02
 FIRENZE DFF 393/08/02
 UGVA-DPT-2002-09/1102

Massive quark effects in two flavor color superconductors

R. Casalbuoni^{a,b}, F. De Fazio^d, R. Gatto^e, G. Nardulli^{c,d} and
 M. Ruggieri^{c,d}

^a*Dipartimento di Fisica, Università di Firenze, I-50125 Firenze, Italia*

^b*I.N.F.N., Sezione di Firenze, I-50125 Firenze, Italia*

^c*Dipartimento di Fisica, Università di Bari, I-70124 Bari, Italia*

^d*I.N.F.N., Sezione di Bari, I-70124 Bari, Italia*

^e*Départ. de Physique Théorique, Université de Genève, CH-1211 Genève 4, Suisse*

Abstract

The high density effective theory formalism (HDET) is employed to describe high density QCD with two massive flavors (2SC). The gap equation is derived and explicitly solved for the gap parameter. The parameters associated to the pseudo Nambu-Goldstone boson of $U(1)_A$ are evaluated in the limit $\mu \rightarrow \infty$ and m/μ fixed. In particular we find for the velocity of the NG boson the relation $v^2 = \sqrt{\mu_1^2 - m_1^2} \sqrt{\mu_2^2 - m_2^2} / 3\mu_1\mu_2$.

1 Introduction

In the present letter we study the effect of quark masses in the two flavor color superconductor [1] (for a recent review see [2]) in a BCS pairing. Correspondingly we will take the masses of two quarks not too different from each other. Massive quarks have been studied by several authors both for two and three flavors [3], [4], [5], [6], [7], [8]. In most of the papers quark masses are treated perturbatively, under the hypothesis that they are much smaller than the energy gap. However this assumption is questionable for the strange quark, which may affect phenomenological analyses (for examples see [9], [10]). In the present paper we make a first attempt to go beyond the above

mentioned approximation by considering the limit $\mu \rightarrow \infty$ with $x \equiv m/\mu$ fixed. We will perform our analysis within the computational scheme of the High Density Effective Theory (HDET), [11, 12] which is most suitable for calculations involving the high density limit of QCD. We will limit most of our analysis to the case of two flavors, though an extension to three flavors is in principle admissible. In Section 2 we establish our notations. Section 3 is dedicated to the 2SC model with massive quarks where we study the gap equation. These results generalize the results of [7]. In Section 4 we analyze the properties of the pseudo Nambu Goldstone Boson (NGB) related to the spontaneous breaking of the $U(1)_A$ symmetry. We recover known results for the mass, although with a different method. On the other hand we find a new mass effect in the velocity of the pseudo NGB, which is given by

$$v^2 = \frac{\sqrt{\mu_1^2 - m_1^2} \sqrt{\mu_2^2 - m_2^2}}{3\mu_1\mu_2}, \quad (1)$$

where μ_k, m_k ($k = 1, 2$) are the masses and quark chemical potentials of the two quarks in the Cooper pair. This result is not in contradiction with the findings of other authors, see e.g. [6], since our corrections are of the order m/μ rather than m/Δ . We also argue that this result should be true at all orders in our parameter expansion.

2 The effective lagrangian for a massive gapless quark

Starting point is the QCD lagrangian

$$\mathcal{L}_{QCD} = \bar{\psi} (i \not{D} + \mu \gamma^0 - m) \psi = \bar{\psi} (i \not{D} + \mu \gamma^0 - x\mu) \psi \quad (2)$$

where we shall held

$$x = \frac{m}{\mu}, \quad 0 < x < 1 \quad (3)$$

fixed in the $\mu \rightarrow \infty$ limit.

We introduce velocity dependent fields $\psi_{\vec{v}}(y)$ and $\Psi_{\vec{v}}(y)$

$$\psi_{\vec{v}}(y) = \mathcal{P}_+ \int_{|\ell| < \delta} \frac{d^4\ell}{(2\pi)^4} e^{i\ell \cdot y} \tilde{\psi}(\ell), \quad \Psi_{\vec{v}}(y) = \mathcal{P}_- \int_{|\ell| < \delta} \frac{d^4\ell}{(2\pi)^4} e^{i\ell \cdot y} \tilde{\psi}(\ell), \quad (4)$$

where $\Delta \ll \delta \ll \mu$ and we have written the quark momentum as $p^\mu = \mu v^\mu + \ell^\mu$, with $v^\mu = (0, \vec{v})$. In (4) we have used the projectors

$$\mathcal{P}_\pm = \frac{1 \pm (\vec{\alpha} \cdot \vec{v} + x \gamma^0)}{2} . \quad (5)$$

These projectors can be obtained from the conventional ones, $(1 \pm \vec{\alpha} \cdot \vec{v})/2$, via a Cini-Touschek transformation [13]. We recall that this transformations was introduced to study the ultra-relativistic limit of the Dirac equation. Owing to (4) we can write

$$\psi(y) = \sum_{\vec{n}} e^{-i\mu \vec{v} \cdot y} [\psi_{\vec{v}}(y) + \Psi_{\vec{v}}(y)] , \quad (6)$$

where the sum represents an average over the velocity directions, with $\vec{n} = \vec{v}/|\vec{v}|$ and arises because there are infinitely many ways to decompose the quark momentum according to (2). $\psi_{\vec{v}}(y)$ and $\Psi_{\vec{v}}(y)$ represent respectively the positive energy and negative energy solution of the Dirac equation associated to the lagrangian (2). We now substitute (6) in $\mathcal{L}_{QCD}$ and make use of the Riemann-Lebesgue lemma to eliminate one of the two sums appearing in the lagrangian (velocity superselection rule of HDET); therefore we get

$$\mathcal{L}_0 = \sum_{\vec{n}} \left\{ \psi_{\vec{v}}^\dagger i \gamma^0 \not{D} \psi_{\vec{v}} + \Psi_{\vec{v}}^\dagger (i \gamma^0 \not{D} + 2\mu) \Psi_{\vec{v}} + \Psi_{\vec{v}}^\dagger i \gamma^0 \not{D} \psi_{\vec{v}} + \psi_{\vec{v}}^\dagger i \gamma^0 \not{D} \Psi_{\vec{v}} \right\} \quad (7)$$

For quarks near the Fermi surface we can write $|\vec{p}| \approx \mu |\vec{v}|$ and $m^2 = \mu^2 (1 - v^2)$; therefore the parameter x defined in (3) is also given by

$$x = \sqrt{1 - v^2} . \quad (8)$$

We will use the identities

$$\begin{aligned} \bar{\psi}_{\vec{v}} \gamma^0 \psi_{\vec{v}} &= \psi_{\vec{v}}^\dagger \psi_{\vec{v}}, & \bar{\psi}_{\vec{v}} \vec{\gamma} \psi_{\vec{v}} &= \psi_{\vec{v}}^\dagger \vec{v} \psi_{\vec{v}} , \\ \bar{\Psi}_{\vec{v}} \gamma^0 \Psi_{\vec{v}} &= \Psi_{\vec{v}}^\dagger \Psi_{\vec{v}}, & \bar{\Psi}_{\vec{v}} \vec{\gamma} \Psi_{\vec{v}} &= -\Psi_{\vec{v}}^\dagger \vec{v} \Psi_{\vec{v}} , \end{aligned} \quad (9)$$

and

$$\mathcal{P}_- \vec{\alpha} \cdot \vec{v} \mathcal{P}_+ = x(x - \gamma_0) P_+ . \quad (10)$$

We can integrate out in the functional integration $\Psi_{\vec{v}}$ through the equation

$$\Psi_{\vec{v}} = \frac{-i \gamma^0 \not{D}}{i \tilde{V} \cdot D + 2\mu} \psi_{\vec{v}} , \quad (11)$$

where we have used the notations $V^\mu = (1, \vec{v})$, $\tilde{V}^\mu = (1, -\vec{v})$

As in the case of massless flavor, the $\Psi_{\vec{v}}$ fields are of order $1/\mu$ with respect to $\psi_{\vec{v}}$ fields; therefore, at the leading order in $1/\mu$ one can write the Dirac part of the massive quark lagrangian as

$$\mathcal{L}_0 = \sum_{\vec{n}} \psi_{\vec{v}}^\dagger i V \cdot D \psi_{\vec{v}} . \quad (12)$$

The integration of the antiquark fields gives a correction to the kinetic term of order x^2/μ . This correction can be neglected in our limit.

Since the sum over velocities is symmetric over the entire solid angle, one can write $\mathcal{L}_0$ as follows:

$$\mathcal{L}_0 = \frac{1}{2} \sum_{\vec{n}} \{ \psi_+^\dagger (i V \cdot D) \psi_+ + \psi_-^\dagger (i \tilde{V} \cdot D) \psi_- \} \quad (13)$$

where the subscripts $\pm$ denote the velocity direction, i.e.

$$\psi_\pm = \psi_{\pm\vec{v}} . \quad (14)$$

This equation is formally the same as in the massless case, however one should remember that the velocity satisfies the condition (8). Another important difference is that we do not separate left and right degrees of freedom, and $\psi_\pm$ have two independent components. Finally the average over the directions is defined as follows:

$$\sum_{\vec{n}} = \int \frac{d\vec{n}}{8\pi} . \quad (15)$$

The extra factor $1/2$ in the average is introduced because, after the introduction of the field with opposite velocity ψ_- , one doubles the degrees of freedom, which implies that the integration must be only over half solid angle. For the following it will be useful to introduce the Nambu-Gorkov notation

$$\chi = \frac{1}{\sqrt{2}} \begin{pmatrix} \psi_+ \\ C\psi_-^* \end{pmatrix} . \quad (16)$$

3 2SC with massive quarks

We consider the model with two massive quarks of different flavor in a color superconductor state [7]. At the Fermi surface one has

$$\vec{p}_1 = \mu_1 \vec{v}_1, \quad \vec{p}_2 = \mu_2 \vec{v}_2 \quad (17)$$

Let us put

$$\mu_1 = \mu - \delta\mu, \quad \mu_2 = \mu + \delta\mu. \quad (18)$$

The Cooper pair in the 2SC model couples quarks of different flavor, and the total momentum of the pair must vanish

$$\vec{p}_1 + \vec{p}_2 = 0. \quad (19)$$

One gets immediately, from $p_1 = p_2$ that

$$\frac{\delta\mu}{\mu} = \frac{\sqrt{1-x_1^2} - \sqrt{1-x_2^2}}{\sqrt{1-x_1^2} + \sqrt{1-x_2^2}} = \frac{m_2^2 - m_1^2}{4\mu^2}, \quad (20)$$

where $x_i = m_i/\mu_i$. Eq. (20) is a consequence of (19) and fixes $\delta\mu$ once the masses and μ are given. Here we are taking advantage of the fact that only strong interactions are considered and therefore $\delta\mu$ has no constraints. By including also weak interactions, this line of reasoning might fail. As discussed in [7], however, if the deviations from (19) and (20) are small, the value of the gap remains inalterd. Only for larger deviations one has a change, since there is a phase transition to the crystalline superconducting state [14], [15]. Eventually, for very large deviations, the normal non superconducting state becomes energetically favorite.

In order to write down the HDET lagrangian we first observe that the Dirac lagrangian not including the gap terms is simply given by two terms (13), one for each flavor. In the 2SC phase we have

$$\langle 0 | \psi_{\alpha i}^T C \gamma_5 \psi_{\beta j} | 0 \rangle = \Delta \epsilon^{\alpha\beta 3} \epsilon_{ij3} \neq 0, \quad (21)$$

where i and j are the flavor indices and α and β are color indices. Therefore we add the term

$$\mathcal{L}_{gap} = \mathcal{L}_{\Delta} + \mathcal{L}_{\bar{\Delta}}, \quad (22)$$

with

$$\mathcal{L}_{\Delta} = - \sum_{\vec{n}} \frac{\Delta}{2} \epsilon^{\alpha\beta 3} \epsilon_{ij3} \psi_{\alpha i, -}^T C \gamma_5 \psi_{\beta j, +} + h.c. \quad (23)$$

and

$$\mathcal{L}_{\bar{\Delta}} = - \frac{\Delta}{2} \epsilon^{\alpha\beta 3} \epsilon_{ij3} \sum_{\vec{n}} \left(\Psi_{\alpha i, -}^T C \gamma_5 \Psi_{\beta j, +} + \psi_{\alpha i, -}^T C \gamma_5 \Psi_{\beta j, +} + \Psi_{\alpha i, -}^T C \gamma_5 \psi_{\beta j, +} \right) + h.c. \quad (24)$$

$\mathcal{L}_{\bar{\Delta}}$ contains antiquark operators. These operators are formally suppressed but, as in the case of the Meissner mass [12], they may give rise to divergent contributions. This happens for the mass of the $U(1)_A$ boson, where we get an unsuppressed term at the leading order in μ . We notice that by the equation (24) we implicitly assume that the antiquark gap $\bar{\Delta}$ is equal to the quark gap Δ . Using eqs. (11) and (10) we can write in momentum space for free quarks

$$\Psi_{\vec{v}} = \frac{-x(x - \gamma_0)\ell_{\parallel}}{|\vec{v}|(2\mu + \tilde{V} \cdot \ell)} \psi_{\vec{v}}, \quad (25)$$

which is proportional to the quark mass.

This introduces in the fermionic lagrangian in the 2SC model the non local interaction term

$$\begin{aligned} \mathcal{L}_{\bar{\Delta}} = & -\frac{\Delta}{2} \epsilon^{\alpha\beta 3} \epsilon_{ij3} \sum_{\vec{n}} \psi_{\alpha i, -}^T C \left\{ \frac{\ell_{\parallel} x_i (x_i + \gamma_0)}{|\vec{v}_i|(2\mu_i + V_i \cdot \ell)} \gamma_5 \frac{\ell_{\parallel} x_j (x_j - \gamma_0)}{|\vec{v}_j|(2\mu_j + \tilde{V}_j \cdot \ell)} \right. \\ & \left. - \gamma_5 \frac{\ell_{\parallel} x_j (x_j - \gamma_0)}{|\vec{v}_j|(2\mu_j + \tilde{V}_j \cdot \ell)} - \gamma_5 \frac{\ell_{\parallel} x_i (x_i - \gamma_0)}{|\vec{v}_i|(2\mu_i + V_i \cdot \ell)} \right\} \psi_{\beta j, +} + h.c. \end{aligned} \quad (26)$$

These terms will be taken in the large μ limit. Since $\mathcal{L}_{\bar{\Delta}}$ can contribute, in the leading order in μ , only to divergent diagrams (i.e. giving rise to μ factors), it may be treated as an insertion.

Let us introduce

$$\chi_{\alpha} = \begin{pmatrix} \chi_{\alpha,1} \\ \chi_{\alpha,2} \end{pmatrix}, \quad (27)$$

where $\chi_{\alpha,i}$ are the Nambu-Gorkov fields (see eq. (16)) for the two flavors. Then we get:

$$\mathcal{L} = \sum_{\vec{n}, \alpha, \beta} \chi_{\alpha}^{\dagger} \begin{pmatrix} iV_1 \cdot D^{\alpha\beta} & 0 & 0 & \hat{\Delta}^{\alpha\beta} \\ 0 & i\tilde{V}_1 \cdot D^{*,\alpha\beta} & \hat{\Delta}^{\alpha\beta} & 0 \\ 0 & -\hat{\Delta}^{\alpha\beta} & iV_2 \cdot D^{\alpha\beta} & 0 \\ -\hat{\Delta}^{\alpha\beta} & 0 & 0 & i\tilde{V}_2 \cdot D^{*,\alpha\beta} \end{pmatrix} \chi_{\beta}, \quad (28)$$

$$\hat{\Delta}^{\alpha\beta} = \gamma_5 \Delta \epsilon^{\alpha\beta 3}. \quad (29)$$

To get the quark propagator we define

$$D_1(\ell, \Delta) = V_1 \cdot \ell \tilde{V}_2 \cdot \ell - \Delta^2, \quad D_2(\ell, \Delta) = V_2 \cdot \ell \tilde{V}_1 \cdot \ell - \Delta^2. \quad (30)$$

We get

$$S^{\alpha\beta}(\ell) = \begin{pmatrix} \frac{\tilde{V}_2 \cdot \ell}{D_1} \delta^{\alpha\beta} & 0 & 0 & -\frac{\hat{\Delta}^{\alpha\beta}}{D_1} \\ 0 & \frac{V_2 \cdot \ell}{D_2} \delta^{\alpha\beta} & -\frac{\hat{\Delta}^{\alpha\beta}}{D_2} & 0 \\ 0 & \frac{\hat{\Delta}^{\alpha\beta}}{D_2} & \frac{\tilde{V}_1 \cdot \ell}{D_2} \delta^{\alpha\beta} & 0 \\ \frac{\hat{\Delta}^{\alpha\beta}}{D_1} & 0 & 0 & \frac{V_1 \cdot \ell}{D_1} \delta^{\alpha\beta} \end{pmatrix}. \quad (31)$$

To get the gap equation in the 2SC model we use the Schwinger-Dyson equation with a four-fermion interaction:

$$\Sigma^{\alpha\beta} = -ig^2 \int \frac{d^4 k}{(2\pi)^4} \frac{g_{\mu\nu} \delta^{ab}}{\Lambda^2} \Gamma_a^{\mu, \alpha\gamma} S_{\gamma\delta}(k) \Gamma_b^{\nu, \delta\beta}, \quad (32)$$

where $\Sigma = -[S^{-1} - S_F^{-1}]$, S_F^{-1} stands for the inverse of the free quark propagator and

$$\Gamma_a^{\mu, \alpha\beta} = \begin{pmatrix} i V_1^\mu \cdot T_a^{\alpha\beta} & & & \\ & -i \tilde{V}_1^\mu \cdot T_a^{T, \alpha\beta} & & \\ & & i V_2^\mu \cdot T_a^{\alpha\beta} & \\ & & & -i \tilde{V}_2^\mu \cdot T_a^{T, \alpha\beta} \end{pmatrix};$$

$T_a^{\alpha\beta}$ being the usual $SU(3)$ generators. We get:

$$\hat{\Delta}^{\alpha\beta} = -i \frac{g^2}{\Lambda^2} \tilde{V}_2 \cdot V_1 (T_a \hat{\Delta} T_a^T)^{\alpha\beta} 4\pi(\mu^2 - \delta\mu^2) \alpha(x_1, x_2) \int \frac{d^2 \ell}{(2\pi)^4} \frac{1}{D_1(\ell, \Delta)} \quad (33)$$

The factor $(\mu^2 - \delta\mu^2) \alpha(x_1, x_2) = \mu_1^2 |\vec{v}_1|^2 = \mu_2^2 |\vec{v}_2|^2 = p_F^2$ implies a reduction of the phase space with respect to the massless case. For the sequel the following notations will be useful:

$$\alpha(x_1, x_2) = \sqrt{1 - x_1^2} \sqrt{1 - x_2^2}, \quad \beta(x_1, x_2) = \sqrt{1 - x_2^2} - \sqrt{1 - x_1^2}. \quad (34)$$

The gap equation, for $\Delta \neq 0$ is

$$1 = i \frac{g^2(\mu^2 - \delta\mu^2) \alpha(x_1, x_2)}{3\Lambda^2 \pi^3} \frac{\tilde{V}_2 \cdot V_1}{2} I(\Delta, x_1, x_2), \quad (35)$$

where

$$I(\Delta, x_1, x_2) = \int \frac{d^2\ell}{\ell_0^2 - \alpha(x_1, x_2)\ell_{\parallel}^2 + \beta(x_1, x_2)\ell_0\ell_{\parallel} - \Delta^2} . \quad (36)$$

where the limits of $\ell_{\parallel}$ integration are $\pm\delta$. One can easily check that for $x_1 = x_2 = 0$ one recovers the gap equation for the 2SC model in the HDET approximation and for massless quarks¹. Performing the integral we obtain:

$$\Delta = \sqrt{\frac{4\alpha + \beta^2}{4}} \frac{\delta}{\sinh\left(\frac{3\Lambda^2\pi^2}{(\mu^2 - \delta\mu^2)\alpha g^2} \frac{\sqrt{4\alpha + \beta^2}}{2(1 + \alpha)}\right)} . \quad (37)$$

If the flavor 1 means the quark up and 2 the strange quark we can write $x_1 \approx 0$; $x \equiv x_2$ and approximate (20) as follows:

$$\frac{\delta\mu}{\mu} = \frac{1 - \sqrt{1 - x^2}}{1 + \sqrt{1 - x^2}} , \quad (38)$$

or

$$\frac{\delta\mu}{\mu} = \frac{m_2^2}{4\mu^2} , \quad (39)$$

and we get

$$\Delta \approx \left(1 - \frac{\delta\mu}{\mu}\right) \frac{\delta}{\sinh\left(\frac{3\Lambda^2\pi^2}{2\mu^2 g^2 (1 - \frac{2\delta\mu}{\mu})}\right)} . \quad (40)$$

In Fig. 1 we show the ratio of the gap for finite masses Δ over the gap for massless quarks Δ_0 as a function of $\delta\mu$. In the plot x_2 has been traded off for $\delta\mu$ and different values of x_1 have been used. Also the difference between the approximate expression and the exact one is given for $x_1 = 0$.

The greatest effect of the strange quark mass on the condensation energy of the BCS pair is due to the reduction of the size of the Fermi surface. This implies that for small values of $\delta\mu$ the condensation energy of the pair decreases linearly with $\delta\mu$. The $x_1 = 0$ curve (dotted line) plots the same quantity as in ref. [7] and agrees quantitatively with the results found there. It can be observed that also the results of [7], similarly to what we have done here, are neither obtained by expanding in m/Δ nor in m/μ . The

¹See eqs. (2.188) and (2.184) of [16].




Fig. 1 The ratio of the gap for finite masses Δ over the gap for massless quarks Δ_0 as a function of $\delta\mu$. The solid line refers to the approximate expression for Δ , see eq. (40), whereas the other three refer to the complete expression in eq. (37) with the dotted line corresponding to $x_1 = 0$, the dash-dotted line to $x_1 = 0.1$, and the dashed one to $x_1 = 0.25$. The values used are $\mu = 400$ MeV, $\Lambda/g = 181$ MeV, corresponding to $\Delta_0 \approx 40$ MeV. Notice that plotting $\delta\mu$ is the same as plotting the squared masses difference, $m_2^2 - m_1^2$ due to the relation (20).

differences between our calculation and that of [7] are as follows. Our eq. (37) presents the advantage to offer an analytical expression; moreover it can be used when both masses, and not only one, are nonvanishing. On the other hand our results are only valid for $m_i \leq \mu$; for higher values the evaluations of [7], that do not suffer of this limitation, should be used.

We note that the insertion of $\mathcal{L}_{\bar{\Delta}}$ would not change the main result we found, mainly the reduction of the phase space, because its effects would only change Δ by factors of order x/μ .

4 $U(1)_A$ pseudo Nambu-Goldstone mode

Besides color symmetry, that is broken from $SU(3)_c$ down to $SU(2)_c$, the condensate (21) also breaks $U(1)_B$ and $U(1)_{em}$; however two linear combinations of these abelian subgroups remain unbroken and there are no NGBs associated to them. The vacuum expectation value (21) breaks spontaneously

$U(1)_A$, which is however also broken by the strong anomaly. At high μ this latter breaking is soft and one expects that the associated pseudo Goldstone boson, the η' , is almost massless. In the sequel we will study the effect on η' of the quark masses in the approximation of neglecting the strong anomaly.

We introduce an external field σ associated to the pseudo Goldstone boson and we put

$$U = \exp \left\{ i \frac{\sigma}{f_\sigma} \right\} .$$

In (16) we perform the substitution

$$\psi_{\alpha_j+} \rightarrow U^\dagger \psi_{\alpha_j+}, \quad \psi_{\alpha_j-}^* \rightarrow U \psi_{\alpha_j-}^*, \quad (41)$$

and $\mathcal{L}_0 + \mathcal{L}_\Delta$ becomes $\mathcal{L}_0 + \mathcal{L}_{\Delta+\sigma}$, with

$$\mathcal{L}_{\Delta+\sigma} = \sum_{\vec{n}} \chi_\alpha^\dagger \hat{\Delta}^{\alpha\beta} \begin{pmatrix} & & U^2 \\ & U^{\dagger 2} & \\ -U^{\dagger 2} & -U^2 & \end{pmatrix} \chi_\beta . \quad (42)$$

At the second order in the σ field:

$$U^2 \approx 1 + \frac{2i\sigma}{f_\sigma} - \frac{2\sigma^2}{f_\sigma^2} \quad (43)$$

and the quark-boson interaction lagrangian can be written as follows:

$$\mathcal{L}_\sigma = \frac{2i\sigma}{f_\sigma} \sum_{\vec{n}} \chi_\alpha^\dagger \hat{\Delta}^{\alpha\beta} \Gamma_0 \chi_\beta - \frac{2\sigma^2}{f_\sigma^2} \sum_{\vec{n}} \chi_\alpha^\dagger \hat{\Delta}^{\alpha\beta} \Gamma_1 \chi_\beta, \quad (44)$$

where

$$\Gamma_0 = \begin{pmatrix} & & 1 \\ & -1 & \\ 1 & -1 & \end{pmatrix}, \quad \Gamma_1 = \begin{pmatrix} & & 1 \\ & 1 & \\ -1 & -1 & \end{pmatrix}. \quad (45)$$

The σ kinetic term arises from loop expansion after functional integration of the fermionic degrees of freedom. Therefore we have the following effective action

$$\mathcal{S}_{eff} = i \frac{1}{2} \text{Tr} \int dx dy \left\{ i S(y, x) \frac{2i\sigma(x)}{f_\sigma} \hat{\Delta} i \Gamma_0 i S(x, y) \frac{2i\sigma(y)}{f_\sigma} \hat{\Delta} i \Gamma_0 \right\}$$

$$+ i\text{Tr} \int dx \left\{ i S(x, x) \frac{(-2)}{f_\sigma^2} \hat{\Delta} \sigma^2(x) i \Gamma_1 \right\} . \quad (46)$$

In evaluating the trace over the spin indices one has to include the $\mathcal{P}_+$ projector which gives a factor 2 if no other γ -matrices are involved. The two terms correspond to the self-energy and a tadpole diagram respectively, see fig. 2.




Fig. 2 One-loop diagrams. External lines represent the pseudo Goldstone field. Full lines are fermion propagators.

The result of the calculation of the effective lagrangian in momentum space is as follows:

$$i \mathcal{L}_{s.e.}(p) = \frac{16 \mu_1 \mu_2 \alpha \Delta^2}{8\pi^3 f_\sigma^2} \sum_{\vec{n}} \int d^2 \ell \left[\frac{V_1 \cdot \ell \sigma \tilde{V}_2 \cdot (\ell + p) \sigma + V_1 \cdot (\ell + p) \sigma \tilde{V}_2 \cdot \ell \sigma - 2 \Delta^2 \sigma^2}{D_1(\ell) D_1(\ell + p)} + 1 \longleftrightarrow 2 \right] \quad (47)$$

$$i \mathcal{L}_{tad} = -\frac{4 \mu_1 \mu_2 \alpha \Delta^2 \sigma^2}{\pi^3 f_\sigma^2} \sum_{\vec{n}} \int d^2 \ell \left(\frac{1}{D_1(\ell, \Delta)} + \frac{1}{D_2(\ell, \Delta)} \right) . \quad (48)$$

We find

$$\mathcal{L}_{s.e.}(p=0) + \mathcal{L}_{tad} = 0 . \quad (49)$$

We have to add the contribution from the antiquark gaps. The pseudo NGB masses in color superconducting phases take contribution from different sources. In the case of the singlet pseudo NGB, as shown in [8], its mass arises from an effective four quark operator obtained by integrating out the electric gluon field; the result is therefore proportional² to $m_1 m_2 \Delta^2 / \mu^2$. In this approach the antiquark gap terms do not appear explicitly as the antiquark field operators are integrated out.

²The calculations in [8] are in the Color Flavor Locking phase and should be adapted to the 2SC case.

Also in the present approach we integrate out the antiquark operators, as discussed above and $\mathcal{L}_{\bar{\Delta}}$ is expressed in terms of the quark fields only. It contributes both to $\mathcal{L}_{s.e.}(0)$ and to $\mathcal{L}_{tad}$, through two additional vertices $\sigma\chi\chi$ and $\sigma\sigma\chi\chi$ that can be constructed analogously to (44) using (26) and (41). We insert these vertices in the expressions for the self-energy and for the tadpole once, since we will take only the leading term in the quark masses to the mass of the Goldstone. The two contributions do not cancel out and one finds

$$i\mathcal{L}_{eff}(0) = \frac{4\Delta^2\alpha\sigma^2}{\pi^3 f_\sigma^2} x_1 x_2 \hat{I} . \quad (50)$$

For $\hat{I}$ we get:

$$\hat{I} = \int_{-\mu}^{+\mu} d^2\ell \frac{(\mu + \ell_{\parallel})^2 \ell_{\parallel}^2}{(2\mu + \tilde{V} \cdot \ell)(2\mu + V \cdot \ell)D(\ell, \Delta)} \approx -2\pi i \mu^2 \log \frac{\mu}{\Delta} + \mathcal{O}(\mu^2) .$$

This is the leading contribution and arises from the first term in (26), since the other two terms are of order μ^2 and not $\mu^2 \log \mu$.

Let us then consider $\mathcal{L}_{eff}(p) - \mathcal{L}_{eff}(0)$.

$$\begin{aligned} i\mathcal{L}_{eff}(p) - i\mathcal{L}_{eff}(0) &= \frac{4\mu_1\mu_2\alpha\Delta^4}{\pi^3 f_\sigma^2} \sum_{\vec{n}} (V_1 \cdot p)\sigma (\tilde{V}_2 \cdot p)\sigma \int \frac{d^2\ell}{[D_1(\ell)]^3} + \\ &+ \frac{2\mu_1\mu_2\alpha\Delta^4}{\pi^3 f_\sigma^2} \sum_{\vec{n}} \int \frac{d^2\ell}{[D_1(\ell)]^3} \times \\ &\times \left([(\tilde{V}_2 \cdot p)\sigma]^2 [V_1 \cdot \ell]^2 + [(V_1 \cdot p)\sigma]^2 (\tilde{V}_2 \cdot \ell)^2 \right) + (1 \leftrightarrow 2) . \end{aligned} \quad (51)$$

One can show that the second term on the r.h.s. of (51) vanishes, while one finds

$$\int \frac{d^2\ell}{[D_1(\ell)]^3} = \frac{-\pi i}{\Delta^4 \sqrt{4\alpha + \beta^2}} . \quad (52)$$

We get therefore

$$\mathcal{L}_{eff}(p) = -\frac{4\mu_1\mu_2\alpha}{\pi^2 f_\sigma^2} \frac{1}{(4\alpha + \beta^2)^{1/2}} \sum_{\vec{n}} (\tilde{V}_2^\mu V_1^\nu + \tilde{V}_1^\mu V_2^\nu) p_\mu \sigma p_\nu \sigma .$$

From which one finally gets, in coordinate space

$$\mathcal{L}_{eff}(\sigma) = \frac{4\mu_1\mu_2\alpha}{\pi^2 f_\sigma^2} \frac{1}{(4\alpha + \beta^2)^{1/2}} \left((\partial_0 \sigma)^2 - \frac{\alpha}{3} (\vec{\partial} \sigma)^2 - m_\sigma^2 \sigma^2 \right) ,$$

where to get the canonical normalization for the kinetic lagrangian one has to put

$$f_\sigma^2 = \frac{8\mu_1\mu_2}{\pi^2} \frac{|\vec{v}_1||\vec{v}_2|}{(|\vec{v}_1| + |\vec{v}_2|)}. \quad (53)$$

Using the result in (50) one finds at the leading order:

$$m_\sigma^2 = 4\Delta^2 \frac{m_1 m_2}{\mu^2} \log \frac{\mu}{\Delta}, \quad (54)$$

which coincides with the result of [6]. It should be noted that in [8] the masses of the NGB bosons in the CFL case do not present any logarithmic enhancement factor. The major difference between our approach and that of [8], a part from the difference in the models that are studied (CFL versus 2SC), is in the coupling between quarks, as we use a BCS four-fermion interaction while in [8] one considers one gluon exchange interaction.

Nonzero quark masses produce a change in the coupling constant f_σ and a reduction of the velocity of the pseudo-NGB in the medium, i.e.

$$v^2 = \frac{|\vec{v}_1||\vec{v}_2|}{3} = \frac{\sqrt{1-x_1^2}\sqrt{1-x_2^2}}{3}, \quad (55)$$

whereas for massless quarks $v^2 = 1/3$. We wish to stress that Eq. (55) constitutes a new result which goes beyond the expansion in m/Δ commonly used in the computation of mass effects in color superconductivity.

Acknowledgements

We wish to thank M. Mannarelli, K. Rajagopal and T. Schäfer for useful discussions. Two of us, R.C. and G. N., wish to thank the DESY theory group, where this work was completed, for the very kind hospitality.

References

- [1] M. G. Alford, K. Rajagopal and F. Wilczek, Phys. Lett. B **422** (1998) 247 [arXiv:hep-ph/9711395].
- [2] K. Rajagopal and F. Wilczek, in Handbook of QCD, M. Shifman ed. (World Scientific 2001), [arXiv:hep-ph/0011333].

- [3] D. T. Son and M. A. Stephanov, Phys. Rev. D **61** (2000) 074012 [arXiv:hep-ph/9910491].
- [4] M. Rho, A. Wirzba and I. Zahed, Phys. Lett. B **473** (2000) 126 [arXiv:hep-ph/9910550].
- [5] D. K. Hong, T. Lee and D. P. Min, Phys. Lett. B **477** (2000) 137 [arXiv:hep-ph/9912531].
- [6] S. R. Beane, P. F. Bedaque and M. J. Savage, Phys. Lett. B **483** (2000) 131 [arXiv:hep-ph/0002209].
- [7] J. Kundu and K. Rajagopal, [arXiv:hep-ph/0112206].
- [8] T. Schafer, Phys. Rev. D **65**, 074006 (2002) [arXiv:hep-ph/0109052].
- [9] P. F. Bedaque and T. Schafer, Nucl. Phys. A **697** (2002) 802 [arXiv:hep-ph/0105150].
- [10] D. B. Kaplan and S. Reddy, Phys. Rev. D **65** (2002) 054042 [arXiv:hep-ph/0107265].
- [11] D. K. Hong, Phys. Lett. **B473** (2000) 118, hep-ph/9812510; D. K. Hong, Nucl. Phys. **B582** (2000) 451, [arXiv:hep-ph/9905523]; S. R. Beane, P. F. Bedaque and M. J. Savage, Phys. Lett. **B483** (2000) 131, [arXiv:hep-ph/0002209].
- [12] R. Casalbuoni, R. Gatto and G. Nardulli, Phys. Lett. B **498** (2001) 179 and Erratum-ibid. B517 (2001) 483, [arXiv:hep-ph/0010321].
- [13] M. Cini and B. Touschek, Nuovo Cimento **7** (1958) 422.
- [14] M. G. Alford, J. A. Bowers and K. Rajagopal, Phys. Rev. D **63** (2001) 074016 [arXiv:hep-ph/0008208].
- [15] J. A. Bowers and K. Rajagopal, Phys. Rev. D **66** (2002) 065002 [arXiv:hep-ph/0204079].
- [16] G. Nardulli, Riv. Nuovo Cimento, **25**, N. 3 (2003), [arXiv:hep-ph/0202037].